

THERE ARE INFINITELY MANY PERRIN PSEUDOPRIMES

JON GRANTHAM

In memory of Daniel Shanks

ABSTRACT. We prove the existence of infinitely many Perrin pseudoprimes, as conjectured by Adams and Shanks in 1982. The theorem proven covers a general class of pseudoprimes based on recurrence sequences. We use ingredients of the proof of the infinitude many Carmichael numbers, along with zero-density estimates for Hecke L-functions.

1. BACKGROUND

In a 1982 paper [1], Adams and Shanks introduced a probable primality test based on third order recurrence sequences. The following is a version of that test.

Consider sequences $A_n = A_n(r, s)$ defined by the following relations: $A_{-1} = s$, $A_0 = 3$, $A_1 = r$, and $A_n = rA_{n-1} - sA_{n-2} + A_{n-3}$. Let $f(x) = x^3 - rx^2 + sx - 1$ be the associated polynomial and Δ its discriminant. (Perrin's sequence is $A_n(0, -1)$.)

Definition. The **signature mod m** of an integer n with respect to the sequence $A_k(r, s)$ is the 6-tuple $(A_{-n-1}, A_{-n}, A_{-n+1}, A_{n-1}, A_n, A_{n+1}) \bmod m$.

Definitions. An integer n is said to have an **S-signature** if its signature mod n is congruent to $(A_{-2}, A_{-1}, A_0, A_0, A_1, A_2)$.

An integer n is said to have a **Q-signature** if its signature mod n is congruent to (A, s, B, B, r, C) , where for some integer a with $f(a) \equiv 0 \pmod n$, $A \equiv a^{-2} + 2a$, $B \equiv -ra^2 + (r^2 - s)a$, and $C \equiv a^2 + 2a^{-1}$.

An integer n is said to have an **I-signature** if its signature mod n is congruent to (r, s, D', D, r, s) , where $D' + D \equiv rs - 3 \pmod n$ and $(D' - D)^2 \equiv \Delta$.

Definition. A **Perrin pseudoprime** with parameters (r, s) is an odd composite n such that either

- 1) $\left(\frac{\Delta}{n}\right) = 1$ and n has an S-signature or an I-signature, or
- 2) $\left(\frac{\Delta}{n}\right) = -1$ and n has a Q-signature.

Adams and Shanks asked if there are infinitely Perrin pseudoprimes. They answered the question, "Almost certainly yes, but we cannot prove it. Almost certainly, there are infinitely many [Carmichael numbers which are Perrin pseudoprimes], and yet it has never been proved that there are infinitely many Carmichael numbers."

In 1994, Alford, Granville and Pomerance [2] proved that there are infinitely many Carmichael numbers. A simple modification of that proof shows that there are infinitely many Lucas and Lehmer pseudoprimes. (If the characteristic polynomial

Date: November 28, 2006.

2000 *Mathematics Subject Classification.* Primary 11Y11; Secondary 11N13 11N25.

is $f(x)$, require all prime divisors p of the Carmichael numbers to have $p \equiv 1 \pmod{4 \operatorname{disc}(f)}$.

In this paper, the techniques of that proof are combined with results about Hecke L-functions to show that there are infinitely many Perrin pseudoprimes. In fact, a more general result can be proven.

Theorem 1.1. *Let $f(x) \in \mathbb{Z}[x]$ be a monic, squarefree polynomial with splitting field K . There are infinitely many Frobenius pseudoprimes with respect to $f(x)$. In fact, there are $\gg N^c$ Carmichael-Frobenius numbers with respect to K which are less than N , for some $c = c(K) > 0$.*

The concept of Frobenius pseudoprime, introduced in [5], provides a generalization of most pseudoprime definitions based on recurrence sequences. A Carmichael-Frobenius number is a pseudoprime with respect to all polynomials with splitting field K . Proving the theorem for the general case allows specialization to other cases.

In particular, we know from [5] that Theorem 1.1 shows that there are infinitely many Perrin pseudoprimes, if we take $f(x) = x^3 - x - 1$. Also from [5], we have that there are infinitely many pseudoprimes in the senses of Gurak [6] and Szekeres [11].

By Proposition 6.1 of [5], in order to prove Theorem 1.1, it suffices to show that there are infinitely many Carmichael numbers n , such that for all $p|n$, $f(x)$ splits completely mod p . The proof will involve modifying the construction in [2] to ensure that each of the prime factors of the Carmichael numbers constructed has this property.

The main result that will be used in this proof is a version of the “prime ideal theorem for arithmetic progressions” that gives a uniform error term, except for a possible exceptional progression.

2. DISTRIBUTION OF PRIMES

Theorems about the distribution of primes in arithmetic progressions are traditionally proved using Dirichlet characters — homomorphisms from the integers mod q to the complex roots of unity. (The map is defined to be zero on integers not coprime to q .) Since we want to prove a theorem about primes in a particular arithmetic progression which split completely, we employ a slightly different sort of Dirichlet character.

We recall the definitions of [9].

Definitions. *Let K be an algebraic number field and \mathfrak{O}_K its ring of integers. A cycle of K is a formal product $\mathfrak{m} = \prod \mathfrak{p}^{m(\mathfrak{p})}$ extending over all of the primes of K , where the $m(\mathfrak{p})$ are nonnegative integers, almost all 0, with $m(\mathfrak{p}) = 0$ for complex \mathfrak{p} and $m(\mathfrak{p}) \leq 1$ for real \mathfrak{p} . Let \mathcal{J} be the group of fractional ideals of \mathfrak{O}_K . Let $\mathcal{J}(\mathfrak{m})$ be the subgroup of \mathcal{J} generated by the finite primes \mathfrak{p} for which $m(\mathfrak{p}) = 0$. Let $P_{\mathfrak{m}}$ be the subgroup of $\mathcal{J}(\mathfrak{m})$ generated by the nonzero ideals of the form $\mathfrak{O}_K \alpha$, where $\alpha \in \mathfrak{O}_K$ is such that $\alpha \equiv 1 \pmod{\mathfrak{p}^{m(\mathfrak{p})}}$ for each finite prime \mathfrak{p} , and $\alpha > 0$ under each embedding of K in the field of real numbers corresponding to a real prime \mathfrak{p} with $m(\mathfrak{p}) = 1$. The norm of a cycle $\mathfrak{m} = \prod \mathfrak{p}^{m(\mathfrak{p})}$ is the number $N(\mathfrak{m}) = \prod N(\mathfrak{p})^{m(\mathfrak{p})}$, where \mathfrak{p} in the second product ranges over only the finite primes, and $N(\mathfrak{p})$ is the norm of \mathfrak{p} in K .*

A **Dirichlet character of K** is a pair consisting of a cycle \mathfrak{m} of K and a group homomorphism $\chi : \mathcal{J}(\mathfrak{m}) \mapsto \mathbb{C}^*$ such that $P_{\mathfrak{m}}$ is contained in the kernel. We call \mathfrak{m} the **modulus** of χ .

Given two Dirichlet characters χ and χ' with moduli \mathfrak{m} and \mathfrak{m}' , we say that χ is **induced** by χ' if $\mathfrak{m}'(\mathfrak{p}) \leq \mathfrak{m}(\mathfrak{p})$ for all \mathfrak{p} and χ is the composition of the inclusion $\mathcal{J}(\mathfrak{m}) \subset \mathcal{J}(\mathfrak{m}')$ with χ' . A Dirichlet character is **primitive** if it is not induced by any character other than itself. The modulus of the unique primitive character inducing a Dirichlet character χ is called the **conductor** of χ .

For a Dirichlet character χ of K , $L(s, \chi)$ is defined to be $\sum \chi(\mathfrak{i})N(\mathfrak{i})^{-s}$, where the sum is over the nonzero ideals of the ring of integers of K and $\text{Re } s > 1$. This sum is absolutely convergent, and $L(s, \chi')$ can be extended to a meromorphic function on the complex plane. It has a simple pole at $s = 1$ if χ' is principal and is holomorphic otherwise.

Let K be the splitting field of f , $n = [K : \mathbb{Q}]$, and $d = \text{disc}(K)$. Let χ be a Dirichlet character mod q (in the traditional sense). We associate to it a Dirichlet character of K in the following way.

Given an ideal $\mathfrak{i} \subset \mathfrak{O}_K$, let $\chi'(\mathfrak{i}) = \chi(N(\mathfrak{i}))$. Then χ' is an example of a Dirichlet character of K with conductor dividing $N(q)$.

Let $\Psi(x, \chi') = \sum_{N(\mathfrak{i}) < x} \chi'(\mathfrak{i})\Lambda(\mathfrak{i})$, where $\Lambda(\mathfrak{i}) = \log N(\mathfrak{p})$ if $\mathfrak{i} = \mathfrak{p}^k$ for some prime ideal \mathfrak{p} , and 0 otherwise. Then

$$\frac{1}{\phi(q)} \sum_{\chi \bmod q} \bar{\chi}(a)\Psi(x, \chi') = \sum_{\substack{N(\mathfrak{i}) < x \\ N(\mathfrak{i}) \equiv a \bmod q}} \Lambda(\mathfrak{i}).$$

We will prove some results about $L(s, \chi')$ that enable us to obtain results about $\Psi(x, \chi')$, and thus about the distribution of primes that split completely in K and lie in a particular residue class.

Lemma 2.1. *Fix a number field K . Let χ be a real Dirichlet character of K mod \mathfrak{m} . Let $M = N(\mathfrak{m})$. Let s be a real number in the range $2 > s > 1$. If χ is principal, then*

$$\frac{L'}{L}(s, \chi) > -\frac{1}{s-1} - c_1 \log 2M,$$

for some $c_1 > 0$, depending on K . If χ is non-principal, and if $L(s, \chi)$ has some real zero $\rho > 0$,

$$\frac{L'}{L}(s, \chi) > \frac{1}{s-\rho} - c_1 \log 2M,$$

and

$$\frac{L'}{L}(s, \chi) > -c_1 \log 2M,$$

if it has no real zero.

Proof. Assume χ is non-principal. From equation (5.9) of [10],

$$\frac{L'}{L}(s, \chi) = B(\chi) + \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho} \right) - \frac{1}{2} \log A(\chi) - \frac{\gamma'_{\chi}(s)}{\gamma_{\chi}},$$

where the sum is over all the non-trivial zeroes of $L(s, \chi)$, $A(\chi) = dM$, $d = \text{disc}(K)$, and $\gamma_\chi(s) = \left[\pi^{-\frac{s+1}{2}} \Gamma\left(\frac{s+1}{2}\right) \right]^b \left[\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \right]^a$ for nonnegative integers a and b depending on χ such that $a + b = n = [K : \mathbb{Q}]$. The exact dependence, described in [10], is irrelevant here.

The $\log A(\chi)$ term can be bounded because $\log A(\chi) = \log dM \ll \log 2M$. $B(\chi)$ is defined implicitly in [10]. By Lemma 5.1 of that paper, we have

$$B(\chi) = -\text{Re} \sum_{\rho} \frac{1}{\rho}.$$

We have from Lemma 5.3 of [10] that

$$\left| \frac{\gamma'_\chi(s)}{\gamma_\chi(s)} \right| \ll n \log(s+2),$$

where the implied constant is absolute.

Thus $\frac{L'}{L}(s, \chi) > \sum_{\rho} \text{Re} \frac{1}{s-\rho} - c_1 \log 2M$, for some $c_1 > 0$. We have that $\text{Re} \frac{1}{s-\rho} = \frac{s - \text{Re} \rho}{|s-\rho|^2} > 0$, so we can omit any part of the sum. We omit anything but one possible real zero.

Thus

$$\frac{L'}{L}(s, \chi) > \frac{1}{s-\rho} - c_1 \log 2M,$$

if $L(s, \chi)$ has a real zero ρ , and

$$\frac{L'}{L}(s, \chi) > -c_1 \log 2M,$$

independent of the existence of real zeros.

Now assume χ is principal. From (5.9) of [10]

$$\frac{L'}{L}(s, \chi) = \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho} \right) - \frac{1}{s} - \frac{1}{s-1} - \frac{1}{2} \log A(\chi) + \frac{\gamma'_\chi(s)}{\gamma_\chi(s)}.$$

By the same arguments as in the non-principal case (and the fact that $\frac{1}{s} < 1$), we have that

$$\frac{L'}{L}(s, \chi) > -\frac{1}{s-1} - c_1 \log 2M.$$

□

The following is a version of the Landau-Page Lemma for Dirichlet L -functions over a number field.

Lemma 2.2. *Given a number field K , there is a computable constant $c_2 > 0$, depending on K , such that for all $T \geq 2$, there is at most one primitive character χ_1 with modulus \mathfrak{m} , $1 \leq N(\mathfrak{m}) < T$ for which $L(s, \chi_1)$ has a zero $\beta_1 + i\gamma_1$ satisfying $\beta_1 \geq 1 - c_2/\log T$ and $|\gamma_1| < T$.*

Proof. We follow the proof in [3], p. 94.

Let χ_1 and χ_2 be primitive characters mod \mathfrak{m}_1 and \mathfrak{m}_2 , respectively, where $N(\mathfrak{m}_1)$ and $N(\mathfrak{m}_2)$ are at most T .

Consider the expression

$$-\frac{L'}{L}(s, \chi_0) - \frac{L'}{L}(s, \chi_1) - \frac{L'}{L}(s, \chi_2) - \frac{L'}{L}(s, \chi_1 \chi_2),$$

where χ_0 is the principal character modulo the gcd of \mathfrak{m}_1 and \mathfrak{m}_2 . (We define $\gcd(\prod \mathfrak{p}^{m_1(\mathfrak{p})}, \prod \mathfrak{p}^{m_2(\mathfrak{p})}) = \prod \mathfrak{p}^{\min(m_1(\mathfrak{p}), m_2(\mathfrak{p}))}$.) This expression is equal to

$$\sum \Lambda(\mathfrak{i})(\chi_0(\mathfrak{i}) + \chi_1(\mathfrak{i}))(\chi_0(\mathfrak{i}) + \chi_2(\mathfrak{i}))N(\mathfrak{i})^{-s} > 0,$$

for $\text{Re } s > 1$.

Assume that $L(s, \chi_1)$ and $L(s, \chi_2)$ have real zeros, β_1 and β_2 respectively. Applying the previous lemma to (1) for real $s > 1$,

$$-\frac{1}{s - \beta_1} - \frac{1}{s - \beta_2} + \frac{1}{s - 1} + c_3 \log T > 0,$$

for some $c_3 > 0$ depending on K , but not T . Rearranging,

$$\frac{1}{s - \beta_1} + \frac{1}{s - \beta_2} < \frac{1}{s - 1} + c_3 \log T.$$

Let $c_2 = \frac{1}{6c_3}$ and assume that each $\beta_i \geq 1 - \frac{c_2}{\log T}$.

Taking $s = 1 + 3c_2/\log T$ gives us $\frac{1}{s - \beta_i} \geq \frac{\log T}{4c_2}$ and $\frac{1}{s - 1} = \frac{\log T}{3c_2}$.

We now have that

$$\frac{\log T}{2c_2} < \frac{\log T}{3c_2} + c_3 \log T.$$

Simplifying, $\frac{1}{6c_2} < c_3$. Substituting the value of c_2 gives the desired contradiction. \square

For each Dirichlet character χ of a field K and real numbers σ, T , in the ranges $\frac{1}{2} \leq \sigma \leq 1$, $T \geq 0$, let $N(\sigma, T, \chi)$ be the number of zeros $s = \beta + i\gamma$ of the Dirichlet L-function $L(s, \chi)$ inside the box $\sigma \leq \beta \leq 1$ and $|\gamma| \leq T$. Let \mathcal{A} be the set of real numbers $A > 2$ for which there exists a number $C_A \geq 1$, such that for all $\sigma \geq 1 - \frac{1}{A}$ and $T \geq 1$,

$$N(\sigma, T, \mathfrak{m}) := \sum_{\chi \bmod \mathfrak{m}} N(\sigma, T, \chi) \leq C_A (N(\mathfrak{m})T^n)^{A(1-\sigma)},$$

for all moduli \mathfrak{m} .

Hilano [8] has shown that every $A \geq 2890$ is in \mathcal{A} . The existence of such an A was first shown by Fogels [4].

Theorem 2.3. *Let K be a number field. For any given $\epsilon > 0$, there exist numbers $x_\epsilon, \eta_\epsilon > 0$, and an integer $q_\epsilon(x)$, all depending on K , such that whenever $x \geq x_\epsilon$ and $x^{1/2} < y < x$,*

$$\left| \sum_{\substack{N(\mathfrak{i}) < y \\ N(\mathfrak{i}) \equiv a \pmod{q}}} \Lambda(\mathfrak{i}) - \frac{y}{\phi(q)} \right| \leq \epsilon \frac{y}{\phi(q)}$$

for all integers q not divisible by $q_\epsilon(x)$, with $(a, q) = 1$ and q in the range $1 \leq q \leq x^{\eta_\epsilon}$. Furthermore $q_\epsilon(x) > \log x$.

Proof. Let $\rho = 3 \log(36C_A/\epsilon)$. Let $\eta_\epsilon = \min(\frac{1}{3An}, \frac{c_2}{n\rho})$. We can require $x_\epsilon > \max(e^{4A\rho/\eta_\epsilon}, 18(C_A/\epsilon)^{2/\eta_\epsilon})$.

We can deduce the following explicit formula from [9], proof of Theorem 3.1: (equations 3.2, 3.3 and the equation following the ‘‘Hence’’ on p. 493).

(1)

$$\sum_{\substack{N(\mathbf{a}) < y \\ N(\mathbf{a}) \equiv a \pmod q}} \Lambda(\mathbf{a}) = \frac{y}{\phi(q)} - \frac{1}{\phi(q)} \sum_{\chi \pmod q} \bar{\chi}(a) \sum_{\substack{L(\beta+i\gamma, \chi)=0 \\ \beta \geq 1/2, |\gamma| \leq T}} \frac{y^{\beta+i\gamma}}{\beta+i\gamma} +$$

$$O\left(n \log y + n \frac{y \log y (\log y + \log dq + \log T)}{T} + n \log dq + ny^{\frac{1}{2}} \log y (\log q + \log y)\right).$$

We have that $\eta_\epsilon < 1/6$, so $\log q < \log y$ and $q < y^{1/3}$. We take $T = x$, so $y < T < y^2$.

The error term in (2) is

$$O\left(ny^{1/2}(\log^2 y + \log y \log d) + n \log d\right).$$

Since d, n are fixed, the error is $O(y^{1/2} \log^2 y) = O(y^{6/7})$, which is less than $\frac{\epsilon}{3} \frac{y}{\phi(q)}$ for y sufficiently large.

The double sum may be bounded by noting that $|y^{\beta+i\gamma}| = y^\beta$, and $\beta + i\gamma \geq \sqrt{1/4 + \gamma^2} \geq (1 + |\gamma|)/3$.

Thus

$$\left| \sum_{\substack{N(\mathbf{a}) < y \\ N(\mathbf{a}) \equiv 1 \pmod q}} \Lambda(\mathbf{a}) - \frac{y}{\phi(q)} \right| \leq \frac{3}{\phi(q)} \sum_{\chi \pmod q} \sum_{\substack{L(\beta+i\gamma, \chi)=0 \\ \beta \geq 1/2, |\gamma| \leq x}} \frac{y^\beta}{1 + |\gamma|} + \frac{\epsilon}{3} \frac{y}{\phi(q)}.$$

Write \sum_σ^α for a sum over all zeros of $\beta + i\gamma$ of $L(s, \chi)$ and over all characters $\chi \pmod q$ where $\sigma \leq \beta < \alpha$ and $|\gamma| < x$. (Each $\beta + i\gamma$ is counted with multiplicity equal to the number of those L-functions for which it is a zero.) To estimate the double sum in (3) we use the upper bounds $y^\beta \leq y^{1-1/(2An)}$ for $\beta \leq 1 - 1/(2An)$, and $y^\beta \leq y$ for $\tau \leq \beta \leq 1$, where $\tau = 1 - \rho/\log x$. In the range $1 - 1/(2An) \leq \beta \leq \tau$, we use the identity $y^\beta = y^{1-1/(2An)} + \log y \int_{1-1/(2An)}^\beta y^\sigma d\sigma$.

Therefore, the double sum in (3) is at most

(2)

$$\sum_{1/2}^\tau \frac{y^{1-1/(2An)}}{1 + |\gamma|} + \log y \sum_{1-1/(2An)}^\tau \frac{1}{1 + |\gamma|} \int_{1-1/(2An)}^\beta y^\sigma d\sigma + y \sum_\tau^1 \frac{y}{1 + |\gamma|}$$

$$\leq y^{1-1/(2An)} \sum_{1/2}^1 \frac{1}{1 + |\gamma|} + \log y \int_{1-1/(2An)}^\tau y^\sigma \left(\sum_\sigma^1 \frac{1}{1 + |\gamma|} \right) d\sigma + y \sum_\tau^1 \frac{1}{1 + |\gamma|}.$$

For $\sigma \geq 1/2$, we have, by partial summation,

$$\sum_\sigma^1 \frac{1}{1 + |\gamma|} \leq N(\sigma, 2, q) + \frac{N(\sigma, x, q)}{x} + \int_2^x \frac{N(\sigma, t, q)}{t^2} dt.$$

By [7], $N(1/2, t, q) < c_4 n q^n t \log qt$. For t in the range $2 \leq t \leq x$, we have $N(1/2, t, q)/t \leq c_4 n q^n \log qx$.

Applying this,

$$\sum_{1/2}^1 \frac{y^{1-\frac{1}{2An}}}{1 + |\gamma|} \leq 2c_4 n q^n y^{1-\frac{1}{2An}} \log qx \left(2 + \int_2^x \frac{dt}{t} \right) \leq 5c_4 n q^n y^{1-\frac{1}{2An}} \log^2 qx.$$

Since we insist that $\eta_\epsilon < \frac{1}{8An^2}$, the first term in (4) is $O(y^{1-1/(3An)})$, which is $< \frac{\epsilon}{18}y$ for y sufficiently large.

If $\sigma \geq 1-1/(2An)$, then $An(1-\sigma) \leq 1/2$, so that for any t in the range $1 \leq t \leq x$, the estimate from [8] shows that $N(\sigma, t, d) \leq C_A q^{An(1-\sigma)} t^{1/2}$. We deduce that

$$\sum_{\sigma} \frac{1}{1+|\gamma|} \leq C_A q^{An(1-\sigma)} \left(3 + \int_2^x \frac{dt}{t^{3/2}} \right) \leq 5C_A q^{An(1-\sigma)}.$$

Using the bound above, the middle term in (4) is

$$(3) \quad \begin{aligned} &\leq 5C_A q^{An} \log y \int_{1-1/(2An)}^{\tau} \left(\frac{y}{q^{An}} \right)^{\sigma} d\sigma \\ &\leq 5C_A q^{An} \frac{\log y}{\log(y/q^{An})} \frac{y}{q^{An}} \left(\frac{y}{q^{An}} \right)^{-(1-\tau)}. \end{aligned}$$

We have that $q^{An} < x^{An\eta} < y^{1/3}$, so $\frac{\log y}{\log(y/q^{An})} < 3/2$. Also, $\left(\frac{y}{q^{An}} \right)^{-(1-\tau)} = \left(\frac{y}{q^{An}} \right)^{-\rho/\log x} < e^{-\frac{2}{3} \log y \rho / \log x} < e^{\frac{1}{3}\rho}$.

Thus the middle term in (4) is

$$\leq 4C_A y e^{-\frac{1}{3}\rho}.$$

And by the way we chose ρ , that is $\leq \frac{\epsilon}{9}y$.

We apply Lemma 2.2 with $T = x^{n\eta_\epsilon}$ and call the exceptional modulus $q_\epsilon(x)$. Then for all moduli less than $x^{n\eta_\epsilon}$ and not divisible by $q_\epsilon(x)$, the L -function has no zeros $\beta + i\gamma$ with $\beta \geq 1 - \rho/\log x$ and $|\gamma| < x^{n\eta_\epsilon}$.

So the third sum above is

$$y \sum_{\tau} \frac{1}{1+|\gamma|} \leq y \frac{N(\tau, x, q)}{x^{n\eta_\epsilon}} \leq C_A y (q^n x^n)^{A(1-\tau)} / x^{n\eta_\epsilon} < C_A y x^{2An\rho/\log x} / x^{n\eta_\epsilon}.$$

This is less than $C_A y x^{-\eta_\epsilon/2}$, by our choice of x . Also by our choice of x , this is less than $C_A y (18C_A/\epsilon)^{2/\eta_\epsilon} x^{-\eta_\epsilon/2} = \epsilon y/9$. Putting these bounds together, we get the desired theorem. \square

Theorem 2.1 of [2] shows, essentially, that the number of primes in an arithmetic progression less than x cannot be too far away from what you expect. Furthermore, it shows this for “most” moduli up to $x^{\frac{5}{12}}$. Our replacement is the following

Theorem 2.4. *Let $f(t) \in \mathbb{Z}[t]$ be a monic polynomial with splitting field K , $[K : \mathbb{Q}] = n$. Then we have real numbers $x_{1/3}, \eta_{1/3} > 0$ and an integer $q_{1/3}(x) > \log x$, depending on K as described in Theorem 2.3, such that the following statement holds. If $q \leq x^{\eta_{1/3}}$, $\gcd(a, q) = 1$, $q_{1/3}(x) \nmid q$, $x \geq x_{1/3}$ and $x^{1/2} < y < x$, then the number of primes $p < y$ that are $a \pmod q$ and such that $f(t)$ splits into linear factors mod p (equivalently, p splits completely in K) is at least $\frac{1}{2\phi(q)n} \pi(x)$.*

Proof. The previous theorem gives that

$$\sum_{\substack{N(\mathfrak{a}) < y \\ N(\mathfrak{a}) \equiv a \pmod q}} \Lambda(\mathfrak{a}) \leq \frac{(2/3)y}{\phi(q)}.$$

The sum contains two types of things we don't want. The first is prime ideal powers. Those can be dispensed of in the usual way. The second is primes that

don't split completely. For primes that do not split completely, we have $N(p) = p^k$, for $k > 1$, so they can be dealt with in the same way as prime powers. We pass to the estimate on the number of primes by standard techniques ([3], p. 112). \square

Henceforth, let $\eta = \eta_{1/3}$ and $q(x) = q_{1/3}(x)$.

3. PRACHAR'S THEOREM

We use the following variant of Prachar's Theorem (c.f. Theorem 3.1 of [2]).

Theorem 3.1. *If L is a squarefree number not divisible by any prime exceeding $x^{\frac{1-\eta}{2}}$ and for which $\sum_{\text{prime } q|L} \frac{1}{q} \leq \frac{1-\eta}{32n}$, then there is a positive integer $k \leq x^{1-\eta}$ with $(k, L) = 1$ such that*

$$\#\{d|L : dk + 1 \leq x, dk + 1 \text{ is prime, splits fully in } K\} \geq \frac{\#\{d|L : 1 \leq d \leq x^\eta\}}{8n \log x}.$$

Proof. Let $\pi_K(x; q)$ denote the number of primes less than x that are 1 mod q and split completely in K .

From Theorem 2.4, we see that for each divisor of L with $1 \leq d \leq x^\eta$ and $(d, q(x)) = 1$,

$$\pi_K(dx^{1-\eta}; d) \geq \frac{\pi(dx^{1-\eta})}{2n\phi(d)} \geq \frac{dx^{1-\eta}}{2n\phi(d) \log x}.$$

Since any prime factor q of L is at most $x^{\frac{1-\eta}{2}}$, we can use Montgomery and Vaughan's explicit version of the Brun-Titchmarsh theorem to get

$$\pi_K(dx^{1-\eta}; dq) \leq \pi(dx^{1-\eta}; dq, 1) \leq \frac{8}{q(1-\eta)} \frac{dx^{1-\eta}}{\phi(d) \log x}.$$

So the number of primes $p \leq dx^{1-\eta}$ with $p \equiv 1 \pmod{d}$ and $(\frac{p-1}{d}, L) = 1$ that split completely is at least

$$\begin{aligned} & \pi_K(dx^{1-\eta}; d) - \sum_{\text{prime } q|L} \pi_K(dx^{1-\eta}; dq) \\ & \geq \left(\frac{1}{2n} - \frac{8}{1-\eta} \sum_{\text{prime } q|L} \frac{1}{q} \right) \frac{dx^{1-\eta}}{\phi(d) \log x} \geq \frac{x^{1-\eta}}{4n \log x}, \end{aligned}$$

for any divisor not divisible by $q(x)$. But at least half of the divisors of L will not be divisible by $q(x)$.

Thus we have at least

$$\frac{x^{1-\eta}}{8n \log x} \#\{d|L : 1 \leq d \leq x^\eta\}$$

pairs (p, d) where $p \leq d^{1-\eta}$ is prime, $p \equiv 1 \pmod{d}$, p splits completely in K , $(\frac{p-1}{d}, L) = 1$, $d|L$ and $1 \leq d \leq x^\eta$. Each such pair (p, d) corresponds to an integer $\frac{p-1}{d} \leq x^{1-\eta}$ which is coprime to L , so there is at least one integer $k \leq x^{1-\eta}$ with $(k, L) = 1$ such that k has at least

$$\frac{1}{8n \log x} \#\{d|L : 1 \leq d \leq x^\eta\}$$

representations as $\frac{p-1}{d}$ with (p, d) as above. Thus for this integer k we have $\#\{d|L : dk + 1 \leq x, dk + 1 \text{ prime, split completely in } K\} \geq \frac{1}{8n \log x} \#\{d|L : 1 \leq d \leq x^\eta\}$. \square

4. INFINITELY MANY FROBENIUS PSEUDOPRIMES

We recall the results from Section 1 of [2].

Theorem 4.1 (Theorem 1.1 of [2]). *If G is a finite abelian group and m is the maximal order of an element in G , then $n(G) < m(1 + \log(\frac{|G|}{m}))$. ($n(G)$ is the length of the longest sequence of (not necessarily distinct) elements of G for which no non-empty subsequence has product the identity.)*

This theorem is due to van Emde Boas and Kruyswijk, and to Meshulam.

Proposition 4.2 (Proposition 1.2 of [2]). *Let G be a finite abelian group, and let $r > t > n = n(G)$ be integers. Then any sequence of r elements of G contains at least $\frac{\binom{r}{t}}{\binom{r}{n}}$ distinct subsequences of length at most t and at least $t - n$, whose product is the identity.*

Theorem 4.3. *Let K be a number field, and let η be the positive real number depending on K defined above. For any $\epsilon > 0$, the number of Carmichael-Frobenius numbers less than x , with respect to a number field K , is at least $x^{\eta/3-\epsilon}$, for sufficiently large x , depending on ϵ and K .*

Proof. Let \mathcal{Q} be the set of primes $q \in (\frac{y^3}{\log y}, y^3]$ for which $q - 1$ is free of prime factors exceeding y . Friedlander has proven that there is a constant $C > 0$ for which

$$|\mathcal{Q}| \geq C \frac{y^3}{\log y}$$

for all sufficiently large y . Let L be the product of the primes $q \in \mathcal{Q}$; then

$$\log L \leq |\mathcal{Q}| \log(y^3) \leq \pi(y^3) \log(y^3) \leq 2y^3,$$

for all large y . Now $\lambda(L)$ is the least common multiple of $\{q - 1\}$ for those primes q that divide L . Since each such $q - 1$ is free of prime factors exceeding y , we know that if the prime power p^a divides $\lambda(L)$ then $p \leq y$ and $p^a \leq y^3$. We let p^{a_p} be the largest power of p with $p^{a_p} \leq y^3$, then

$$\lambda(L) \leq \prod_{p \leq y} p^{a_p} \leq \prod_{p \leq y} y^3 = y^{3\pi(y)} \leq e^{6y}$$

for all large y .

Let G be the group $(\mathbb{Z}/L\mathbb{Z})^*$. From Theorem 1.1 of [2] and the above equations,

$$n(G) < \lambda(L) \left(1 + \log \frac{\phi(L)}{\lambda(L)} \right) \leq \lambda(L)(1 + \log L) \leq e^{9y}$$

for all large y .

We can choose y large enough so that $\sum \frac{1}{q} \leq \frac{1-B}{32n}$ as needed to apply Theorem 3.1. Let $\delta = \frac{3\epsilon}{8n\eta}$, and let $x = e^{y^{1+\delta}}$. Then, for y large enough, there is an integer k coprime to L for which the set \mathcal{P} of primes $p \leq x$ with $p = dk + 1$ for some divisor d of L , and that split in K , satisfies

$$|\mathcal{P}| \geq \frac{\#\{d|L : 1 \leq d \leq x^\eta\}}{8n \log x}.$$

The product of any

$$u := \left[\frac{\log(x^\eta)}{\log y^3} \right] = \left[\frac{\eta \log x}{\theta \log y} \right]$$

distinct prime factors of L is a divisor d of L with $d \leq x^\eta$. We deduce from above that

$$\begin{aligned} \#\{d|L : 1 \leq d \leq x^\eta\} &\geq \binom{\omega(L)}{u} \geq \left(\frac{\omega(L)}{u}\right)^u \\ &\geq \left(\frac{Cy^3}{\eta \log x}\right)^u = \left(\frac{C}{\eta}y^{2-\delta}\right)^u. \end{aligned}$$

We notice that $\frac{(2-\delta)\eta}{3} = \frac{2\eta}{3} - \frac{\epsilon}{4}$. So for all sufficiently large values of y ,

$$|\mathcal{P}| \geq \frac{\left(\frac{C}{\eta}y^{2-\delta}\right)^u}{8n \log x} \geq x^{\frac{2\eta}{3} - \frac{\epsilon}{3}}.$$

Take $\mathcal{P}' = \mathcal{P} \setminus \mathcal{Q}$. Since $|\mathcal{Q}| \leq y^3$, we have that $|\mathcal{P}'| \geq x^{\frac{2\eta}{3} - \frac{\epsilon}{2}}$, for all sufficiently large values of y .

We may view \mathcal{P}' as a subset of the group $G = (\mathbb{Z}/L\mathbb{Z})^*$ by considering the residue class of each $p \in \mathcal{P}' \pmod L$. If \mathcal{S} is a subset of \mathcal{P}' that contains more than one element and if

$$\prod(\mathcal{S}) := \prod_{p \in \mathcal{S}} p \equiv 1 \pmod L,$$

then $\prod(\mathcal{S})$ is a Carmichael number all of whose prime factors split completely in K . Thus it is a Frobenius pseudoprime.

Let $t = e^{y^{\frac{1+\delta}{2}}}$. Then, by Proposition 1.2 of [2], we see that the number of Frobenius pseudoprimes of the form $\prod(\mathcal{S})$, where $\mathcal{S} \subset \mathcal{P}'$ and $|\mathcal{S}| \leq t$, is at least

$$\frac{\binom{|\mathcal{P}'|}{t}}{\binom{|\mathcal{P}'|}{n(G)}} \geq \frac{\binom{|\mathcal{P}'|}{t}^{[t]}}{|\mathcal{P}'|^{n(G)}} \geq \left(x^{\frac{2\eta}{3} - \frac{\epsilon}{2}}\right)^{[t] - n(G)} [t]^{-[t]} \geq x^{t(\frac{2\eta}{3} - \epsilon)}$$

for all sufficiently large values of y . We note that we have formed each Frobenius pseudoprime $\prod(\mathcal{P}) \leq x^t$. Thus for $X = x^t$ we have the number of Frobenius pseudoprimes $\leq x$ is at least $X^{\frac{2\eta}{3} - \epsilon}$ for all sufficiently large values of X . Since y can be uniquely determined from X , the theorem is proven. \square

REFERENCES

- [1] W. W. Adams and D. Shanks. Strong primality tests that are not sufficient. *Math. Comp.*, 39:255–300, 1982.
- [2] W. R. Alford, Andrew Granville, and Carl Pomerance. There are infinitely many carmichael numbers. *Annals of Mathematics*, 140:703–722, 1994.
- [3] Harold Davenport. *Multiplicative Number Theory*. Springer-Verlag, New York, second edition edition, 1980.
- [4] E. Fogels. On the zeros of l -functions. *Acta Arith.*, 11:67–96, 1965.
- [5] J. Grantham. Frobenius pseudoprimes. *Math. Comp.*, 70:873–891, 2001.
- [6] S. Gurak. Pseudoprimes for higher-order linear recurrence sequences. *Math. Comp.*, 55:783–813, 1990.
- [7] Teluhiko Hilano. On the zeros of hecke’s l -functions. *Sci. Papers College Gen. Ed. Univ. Tokyo*, 24:8–24, 1974.
- [8] Teluhiko Hilano. On the zeros of heck’s [sic] l -functions. *Proc. Japan Acad.*, 50:23–28, 1974.
- [9] H.W. Lenstra Jr. and Carl Pomerance. A rigorous time bound for factoring numbers. *J. Amer. Math. Soc.*, 5:483–516, 1992.
- [10] J.C. Lagarias and A.M. Odlyzko. Effective versions of the chebotarev density theorem. In A. Frolich, editor, *Algebraic Number Fields: L-functions and Galois Properties*, pages 409–464. Academic Press, London, 1977.

- [11] G. Szekeres. Higher order pseudoprimes in primality testing. In *Combinatorics, Paul Erdős is eighty*, volume 2, pages 451–458. János Bolyai Math Soc., Budapest, 1996.

INSTITUTE FOR DEFENSE ANALYSES, CENTER FOR COMPUTING SCIENCES, 17100 SCIENCE DRIVE,
BOWIE, MD 20715

E-mail address: `grantham@super.org`